

## MISSION:

Take control of infestation


**R&C™** 2in1 Shampoo + Conditioner

### Help eliminate head and pubic (crab) lice using R&C™ Shampoo/Conditioner

Please read and follow treatment instructions **CAREFULLY**.

**Do not** use R&C™ Shampoo/Conditioner on wet hair. **Apply on DRY** hair only.

*Shake product well before using.*



With scissors snip off the tip of the applicator cap.



Unscrew cap on bottle and replace with applicator cap.



**Alert!**  
Do not wet hair before application

Apply R&C™ Shampoo/Conditioner to dry hair.

Apply enough R&C™ Shampoo/Conditioner to thoroughly soak the hair and skin of the infested area and adjacent hairy areas (min 25 mL). Ensure that the shampoo/conditioner is applied close to the scalp for head lice, or close to the skin of the infested area for crab lice. **Be careful to thoroughly soak the nape of the neck and the back of the ears.**



Allow diluted R&C™ Shampoo/Conditioner to remain on the area for 10 minutes.



Add small quantities of water working R&C™ Shampoo/Conditioner into the hair and skin until a lather forms.

R&C™ Shampoo/Conditioner will only kill lice effectively if used on dry hair and as directed.

**Alert!** A second application is usually recommended 7-10 days after the first application if you see live lice or nits (eggs).



Rinse thoroughly.



**Alert!**  
Lice may take up to 24 hours to die after treatment

Remove dead lice and nits with the in-pack nit comb using the following steps:

- Comb wet hair with a regular comb to remove snarls
- Separate a strand of hair not wider than the nit comb
- Using the nit comb tilted toward the scalp, comb the strand from the scalp to the end of the hair
- Wipe off the comb and repeat until all the hair has been combed
- Check the areas behind the ears and around the nape of the neck




Store unused product by removing the applicator cap from the bottle and replacing it with the original cap. Store the product between 15-30° C. Rinse the applicator cap and air dry.

**9** A second application is usually recommended 7-10 days after the first application if you see live lice or nits (eggs).

## MISSION:

Take control of infestation

### Understanding your enemy



A complete louse life cycle from egg, to adult, to egg may take as little as 17 days. Lice may take up to 24 hours to die after initial treatment. A second application of R&C™ Shampoo/Conditioner is recommended 7-10 days after initial application to break the life cycle, serving to kill newly hatched nymphs before they can lay eggs. Following the treatment instructions carefully is necessary to ensure effective treatment of lice infestation and to minimize the risk of re-infestation.<sup>2</sup>

**Alert!** Remember, re-treatment 7-10 days after initial treatment is recommended to ensure lice and nymphs are killed effectively<sup>1,2</sup>

Please consult package insert for cautions and first aid instructions.

Additional instructions:

Enter Treatment Date:	Enter Re-treatment Date:
DD/MM	DD/MM
	7-10 days after first treatment

For more information on how to take control of head lice visit [www.liceguide.com](http://www.liceguide.com)

1. R&C™ Labelling, 2. Canadian Pharmaceutical Association. Self-Medication Reference for Professionals Volume One, 4th Ed. Canadian Pharmaceutical Association, Ottawa, Canada 1992. p 267-277.

### Protect against re-infestation

To prevent head lice from coming back follow these instructions carefully:

- Notify any person who has been in close contact with you so that other cases can be found and treated in the same time frame
- Soak brushes and combs in hot water above 54° C or in undiluted R&C™ Shampoo/Conditioner for 5-10 minutes then rinse thoroughly with hot water
- Articles such as clothing, towels and bed linens may harbour lice or nits. To clean:
  - Wash all clothing, towels, bedding, etc. in very hot water and dry in the dryer by using the hot cycle for at least 20 minutes, **or**
  - Seal in a plastic bag for two weeks and wash normally

Available in the following sizes: 50 mL and 200 mL



Should not be used by persons allergic to chrysanthemum, pyrethrin, or synthetic pyrethroids. Do not use on eyelashes or eyebrows. Do not exceed 2 applications in 24 hours. For external use only. Harmful if swallowed. Do not inhale. Contact should be avoided with eyelashes, eyes, nose and mouth, and other mucosal areas. In the case of infection or skin irritation, discontinue use and consult a physician.

**gsk**  
GlaxoSmithKline  
Consumer Healthcare

©Reg'd TM of GlaxoSmithKline  
GlaxoSmithKline Consumer Healthcare Inc.  
Oakville, Ontario L6H 5V2  
©2004 GlaxoSmithKline

PAAB